



TITLE:

Production of polyhydroxybutyrate and alginate from glycerol by *Azotobacter vinelandii* under nitrogen-free conditions

AUTHOR(S):

Yoneyama, Fuminori; Yamamoto, Mayumi;
Hashimoto, Wataru; Murata, Kousaku

CITATION:

Yoneyama, Fuminori ...[et al]. Production of polyhydroxybutyrate and alginate from glycerol by *Azotobacter vinelandii* under nitrogen-free conditions. *Bioengineered* 2015, 6(4): 209-217

ISSUE DATE:

2015-04-16

URL:

<http://hdl.handle.net/2433/207605>

RIGHT:

This is an Accepted Manuscript of an article published by Taylor & Francis in 'Bioengineered' on 16 Apr 2015, available online: <http://www.tandfonline.com/10.1080/21655979.2015.1040209>; The full-text file will be made open to the public on 16 April 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。 ; This is not the published version. Please cite only the published version.

1 *MS: for Bioengineered as a Research article*

2
3 **Title:** Production of polyhydroxybutyrate and alginate from glycerol by
4 *Azotobacter vinelandii* under nitrogen-free conditions

5
6 **Running title:** Production of biopolymers from glycerol

7
8 **Authors:** Fuminori Yoneyama¹, Mayumi Yamamoto¹, Wataru Hashimoto¹,
9 Kousaku Murata^{1,2*}

10
11 **Affiliations:** ¹*Division of Food Science and Biotechnology, Graduate School of*
12 *Agriculture, Kyoto University, Uji, Kyoto 611-0011, Japan*

13 ²*Present address: Department of Life Science, Faculty of Science and*
14 *Engineering, Setsunan University, Neyagawa, Osaka 572-8508, Japan*

15
16 *To whom correspondence should be addressed.

17 Department of Life Science, Faculty of Science and Engineering, Setsunan University,
18 Neyagawa, Osaka 572-8508, Japan. Phone: +81 72 800 1036. Fax: +81 72 800 1036.
19 E-mail: k-murata@lif.setsunan.ac.jp.

20
21 **Keywords:** alginate, azotobacter, glycerol, nitrogen fixation, polyhydroxybutyrate

22
23 **Competing financial interests:** The authors declare no competing financial interests.

ABSTRACT

Glycerol is an interesting feedstock for biomaterials such as biofuels and bioplastics because of its abundance as a by-product during biodiesel production. Here we demonstrate glycerol metabolism in the nitrogen-fixing species *Azotobacter vinelandii* through metabolomics and nitrogen-free bacterial production of biopolymers, such as poly-D-3-hydroxybutyrate (PHB) and alginate, from glycerol. Glycerol-3-phosphate was accumulated in *A. vinelandii* cells grown on glycerol to the exponential phase, and its level drastically decreased in the cells grown to the stationary growth phase. *A. vinelandii* also overexpressed the glycerol-3-phosphate dehydrogenase gene when it was grown on glycerol. These results indicate that glycerol was first converted to glycerol-3-phosphate by glycerol kinase. Other molecules with industrial interests, such as lactic acid and amino acids including γ -aminobutyric acid, have also been accumulated in the bacterial cells grown on glycerol. Transmission electron microscopy revealed that glycerol-grown *A. vinelandii* stored PHB within the cells. The PHB production level reached 33% per dry cell weight in nitrogen-free glycerol medium. When grown on glycerol, alginate-overproducing mutants generated through chemical mutagenesis produced two-fold the amount of alginate from glycerol than the parental wild-type strain. To the best of our knowledge, this is the first report on bacterial production of biopolymers from glycerol without addition of any nitrogen source.

INTRODUCTION

Azotobacter vinelandii is a free living, nitrogen-fixing bacterium.^{1,2} One of the remarkable characteristics of this species is that it can grow sufficiently in a nitrogen-free minimal medium.³ In *A. vinelandii*, nitrogenases prerequisite for nitrogen fixation catalyze the reduction of nitrogen to ammonia using a large amount of energy derived from ATP hydrolysis.^{4,5} In the recently determined genome sequence of *A. vinelandii*, three different types of nitrogenases have been identified.² In addition, it has been established that *A. vinelandii* has the potential to produce industrially useful biopolymers, including extracellular alginate and intracellular poly-D-3-hydroxybutyrate (PHB).⁶⁻⁹ Alginate is a linear polysaccharide consisting of (1-4)- β -D-mannuronic acid and α -L-guluronic acid. Commercially available alginate is classically derived from seaweed, although two bacterial genera, *Pseudomonas* and *Azotobacter*, are expected to be potential alginate producers.⁸ PHB belongs to the polyhydroxyalkanoate (PHA) family of polyesters, and many bacterial species accumulate PHAs as intracellular granules for energy storage.¹⁰ PHAs are also promising alternatives to plastics because of their biodegradability, biocompatibility, and thermoplasticity.^{6,11} Hence, *A. vinelandii* is considered to be an attractive bacterium for production of two industrially useful biopolymers, alginate and PHB, in the absence of nitrogen sources.^{6,9,12-14}

The biosynthetic pathways for alginate and PHB production in *A. vinelandii* have previously been reviewed by Galindo *et al.*⁶ As shown in Fig. 1, alginate is synthesized from fructose 6-phosphate by many enzymes encoded by the *alg* cluster,^{8,15} whereas PHB is synthesized in three steps from acetyl-CoA and three *phb* genes are essential for synthesis.¹⁶ The regulatory mechanisms for the production of these biopolymers have been analyzed.¹⁶⁻²⁵ Thus, the biopolymers are expected to be produced from excess

and/or unused resource by *A. vinelandii*.

A. vinelandii is known to assimilate various carbon sources. Sucrose, glucose, fructose, mannose, sorbitol, mannitol, glycerol, gluconate, and acetate can all be used as the sole carbon source for cell growth.^{26,27} Furthermore, some of them are also used for biopolymer production.²⁶ However, little information exists on bacterial glycerol metabolism including biopolymer production in a nitrogen-free environment. Glycerol is generated as a major by-product during biodiesel production, and its efficient utilization is now sought in various areas of food, pharmaceutical, agricultural, and environmental research. Moreover, microorganism-mediated conversion of glycerol to other valuable materials is being developed worldwide. Production of hydrogen, ethanol, butanol, 1,3-propanediol, propionic acid, and PHAs has previously been undertaken using bacteria.^{28,29}

To the best of our knowledge, no report exists on glycerol utilization in a nitrogen-free environment. Moreover, glycerol metabolism and production of biopolymers from glycerol in *A. vinelandii* remain to be clarified. The present study involves metabolomics-based identification of the glycerol metabolic pathway in *A. vinelandii* and the bacterial production of biopolymers from glycerol in a nitrogen-free environment.

RESULTS AND DISCUSSION

Sucrose and glycerol metabolism. Metabolites in *A. vinelandii* cells grown on sucrose or glycerol were analyzed by a capillary electrophoresis time-of-flight mass spectrometry (CE-TOFMS) (Tables 1 and 2). Small amounts of glucose-, fructose-, and glycerol-related metabolites, but not glycerol-3-phosphate accumulated in the cells

(Table 1). Considerable amounts (400 pmol/OD₆₀₀ ml) of glycerol-3-phosphate were detected in the glycerol-grown cells collected at the exponential growth phase, whereas its level drastically decreased (82 pmol/OD₆₀₀ ml) in cells collected at the stationary growth phase. Moreover, the dihydroxyacetone phosphate level in the glycerol-grown cells was much higher than that in the sucrose-grown cells. These results suggest that glycerol was first catabolized to glycerol-3-phosphate, then to dihydroxyacetone phosphate, and finally converged into the glycolytic pathway. Because cell growth on glycerol was slower than that on sucrose as described later, accumulation of glycerol-3-phosphate was considered to be a rate-limiting step.

In the recently determined genome sequence of *A. vinelandii* strain DJ,² putative genes for glycerol uptake and degradation were found at a locus containing four genes, *glpF*, *glpK*, *glpR*, and *glpD*. GlpF and GlpR are annotated as putative glycerol-uptake and -repressor proteins, respectively, whereas GlpK and GlpD are annotated as glycerol kinase and glycerol-3-phosphate dehydrogenase, respectively. Although GlpK and GlpD are classified into a group involved in phospholipid metabolism according to genome annotation, metabolomics in the present study demonstrated that glycerol and glycerol-3-phosphate were substrates of GlpK and GlpD, respectively, and that both enzymes were necessary for glycerol metabolism.

Two pathways were postulated for the conversion of glycerol to dihydroxyacetone phosphate. One is through glycerol-3-phosphate and the other through dihydroxyacetone. A number of microorganisms can use glycerol as the sole carbon source through dihydroxyacetone.²⁹⁻³¹ Furthermore, the genome sequence of *A. vinelandii* revealed the presence of glycerol dehydrogenase- and dihydroxyacetone kinase-like genes (Fig. 1). Hence, a dehydrogenase for glycerol and one for glycerol-3-phosphate were assayed to

determine the main glycerol assimilation pathway. However, no activity of either enzyme was detected in the bacterial cell extract, possibly due to their low protein expression level. Hence, transcription levels of the genes of the two dehydrogenases were monitored by quantitative PCR (Fig. 2). As shown in Fig. 2A, the total RNA of bacteria grown in all conditions was extracted without degradation. In the case of the glycerol-3-phosphate dehydrogenase gene (Fig. 2B left), the cells grown to the exponential growth phase showed higher gene expression than those grown to the stationary growth phase. In bacteria grown to the exponential growth phase, gene expression in the glycerol medium was 38-fold higher than that of bacteria grown in the sucrose medium. In contrast, the glycerol dehydrogenase gene was transcribed at basal levels in bacteria grown to the exponential growth phase in media (Fig. 2B right). The expression of glycerol dehydrogenase in all cases tested was extremely low, but near the detectable limit for our experimental conditions (data not shown). These metabolomic and quantitative PCR results demonstrate that the glycerol-3-phosphate pathway is predominant in *A. vinelandii*.

Sucrose metabolites, such as fructose 6-phosphate, mannose 6-phosphate, mannose 1-phosphate, and GDP mannose, were detected even in glycerol-grown *A. vinelandii* (Table 1). This result indicates that gluconeogenesis, which involves the conversion of fructose 1,6-biphosphate to fructose 6-phosphate, occurred in the bacteria. In addition, occurrence of gluconeogenesis in *A. vinelandii* demonstrates its potential for producing alginate concomitantly with PHB from glycerol, which is described later.

No conspicuous rate-limiting metabolites were identified between fructose 1,6-biphosphate and acetyl CoA (Table 1). However, accumulation of pyruvic acid (42.5 pmol/OD₆₀₀ ml) was observed in glycerol-grown cells collected at the exponential

growth phase. Pyruvic acid is an important precursor for many metabolites, including ethanol, acetyl CoA, and lactic acid.³² In the present study, *A. vinelandii* produced large amounts of lactic acid in all cases, especially in glycerol-grown cells (Table 2), and levels of several amino acids accumulated in glycerol-grown cells were higher than those in sucrose-grown cells. Both glutamine and glutamic acid were considerably produced in all cases, especially in glycerol-grown cells collected at the exponential growth phase. Arginine, lysine, and proline are synthesized from glutamic acid in most bacteria. A similar accumulation profile of these amino acids was observed in *A. vinelandii*, as shown in Table 2. Thus, *A. vinelandii* demonstrated the potential for amino acid production under nitrogen-free conditions. Furthermore, γ -aminobutyric acid (GABA), a neurotransmitter of clinical interest,³³ was stored in the cells grown to the exponential growth phase in sucrose and glycerol. Because bacteria-produced GABA is known as a molecule of intracellular pH management due to decarboxylation of glutamic acid,³⁴ this molecule is suggested to play an important role in the neutralization of organic acids accumulated in the cells.

Metabolomic analysis performed in the present study supported the role of the metabolic pathway predicted previously, as shown in Fig. 1. In addition, based on results of the present study, *A. vinelandii* is promising as a potential producer of many useful materials, such as amino acids, GABA, lactic acid, and biopolymers, in a nitrogen-free environment.

PHB production. PHB accumulation was investigated in wild-type (WT) *A. vinelandii* grown on sucrose or glycerol. In addition, the mutant $\Delta algD$ cells, which have a disrupted alginate synthetic gene *algD*, were also subjected to the PHB assay because

PHB and alginate productions are thought to be competitive. First, intracellular PHB granules were monitored by transmission electron microscopy (TEM) (Fig. 3). Similar to many PHB-producing bacteria, WT cells produced white and globular PHB granules, which were found in both sucrose- and glycerol-grown cells. On the other hand, globular PHB granules were scarcely observed and apparently degraded in *ΔalgD* cells.

The time course of cell growth and PHB production are shown in Fig. 4. In the sucrose medium, WT and *ΔalgD* cells showed similar growth profiles (Fig. 4A). Cell growth of both strains exceeded an OD₆₀₀ of 14. However, the lag phase of both strains grown on glycerol was longer than that on sucrose. WT growth reached an OD₆₀₀ of 6.12, whereas that of *ΔalgD* reached an OD₆₀₀ of 16.8.

Figure 4B indicates the time course of intracellular PHB accumulation (per dry cell). The intracellular PHB level in WT cells grown on sucrose as well as glycerol reached approximately 33%. Interestingly, the PHB level of *ΔalgD* cells grown on either source initially increased, but subsequently decreased. As shown in Fig. 4A, PHB was degraded in *ΔalgD* cells. *A. vinelandii* is known to convert from vegetative cells to cysts under unfavorable environments for growth, and PHB has been observed in the cyst-forming cells as a probable energy and carbon storage material.¹⁴ Although the reasons for PHB degradation have yet to be determined, the necessity for PHB might waver due to the lack of alginate production following cyst formation. Production of PHB and alginate is closely regulated in *A. vinelandii*.¹⁰ The results obtained herein may provide valuable hints on the relationships among alginate, PHB, and cyst formation.

Although *A. vinelandii* grown on glucose together with nitrogen sources synthesizes much PHB (74% per dry cell),³⁵ and some bacteria such as *Burkholderia cepacia*,³⁶ *Chelatococcus daeguensis*,³⁷ *Cupriavidus necator* (formerly *Ralstonia eutropha*),^{38,39}

190 *Paracoccus denitrificans*,⁴⁰ *Pseudomonas oleovorans*,⁴¹ and *Zobellella denitrificans*,⁴²
191 have also been known to produce PHB from glycerol with nitrogen sources, the
192 bacterial production of PHB from glycerol without a nitrogen source contributes to
193 green chemistry.

194

195 **Alginate production.** The alginate production level was determined in *A. vinelandii*
196 grown on sucrose or glycerol. The sucrose-grown cells obtained at the stationary phase
197 produced alginate at 0.3–0.4 mg/ml, whereas those grown in the glycerol medium
198 produced enhanced levels of approximately 0.5 mg/ml. To elevate the alginate
199 production level, a random NTG-treated mutation library was used to screen
200 alginate-overproducing mutants. Unlike for PHB, it was easy to select
201 alginate-overproducing mutants because these mutants seemed to form high mucoid
202 colonies. More than 100 mutants were isolated as high mucoid colonies in comparison
203 with WT colonies.

204 One of the mutants (MT1) was subjected to TEM analysis after preparation of cell
205 thin section (Fig. 5A). Recently, *A. vinelandii* cells were demonstrated to be equipped
206 with a special secretion system for alginate through formation of cell-surface blebs.
207 These blebs containing alginate are formed on the bacterial cell surface and are
208 subsequently released around the cells.⁴³ MT1 cells formed a large number of blebs on
209 the cell surface (Fig. 5A). On the other hand, few blebs were observed on WT cells (Fig.
210 5B). Thus, WT cells were subjected to scanning electron microscopy (SEM) analysis. A
211 few blebs were observed on the cell surface (Fig. 5C). The time course of alginate
212 secretion for WT and MT1 cells grown on glycerol is characterized in Fig. 6. Cell
213 growth (3.97 at OD₆₀₀) of MT1 at 96 h was lower than that (6.64 at OD₆₀₀) of WT (Fig.

6A). In contrast, alginate secretion by MT1 cells (0.87 mg/ml) was higher than that by WT cells (0.52 mg/ml) (Fig. 6B). This result indicates that MT1 cells have the potential for more alginate production through improvement in their growth conditions.

In conclusion, this is the first report on glycerol metabolism in *A. vinelandii* analyzed through the metabolomic approach. The results indicate that *A. vinelandii* grown on glycerol, but in the absence of any nitrogen source, may be useful for producing many substances, including amino acids and biopolymers (especially PHB and alginate), with industrial interests.

MATERIALS AND METHODS

Bacteria and culture conditions. The bacterial strains used in the present study are listed in Table 3. Cells of WT *A. vinelandii* ATCC 12837 and an alginate-deficient mutant with a disruption of *algD* ($\Delta algD$)⁴³ were grown aerobically in a minimal glycerol medium, i.e., modified Burk's medium (G-MB; 20 mg/ml glycerol, 200 µg/ml NaCl, 50 µg/ml CaSO₄, 200 µg/ml MgSO₄ 7H₂O, 2.9 µg/ml Na₂MoO₄ 2H₂O, 27 µg/ml FeCl₃, 0.66 mg/ml K₂HPO₄, and 0.16 mg/ml KH₂PO₄) at 30°C with agitation of 120 strokes per min. Sucrose (20 mg/ml) was also used for a carbon source instead of glycerol in G-MB (S-MB).

Metabolome analysis. Metabolome analysis was supported by Human Metabolome Technologies (Tsuruoka, Japan). Precultured WT cells were inoculated in 50 ml fresh S-MB or G-MB, and grown to the exponential or stationary growth phase. To reach the stationary growth phase, cells were grown in each medium for 84 h. To reach the exponential growth phase, cells were grown in S-MB and G-MB for 40 and 60 h

(approximately 1.2 at OD₆₀₀), respectively. The cells were harvested by centrifugation (5700 ×g, 4°C, 5 min), and were washed twice with 10 ml pure water. Each cell pellet was homogenized in 2 ml methanol. After homogenization, chloroform (1.6 ml) was added to 1.6 ml of cell extract, and the mixture was well agitated. Aqueous and chloroform layers were separated by centrifugation (2300 ×g, 4°C, 5 min), and the aqueous layer was subjected to ultrafiltration (9100 ×g, 4°C, 120 min) using the Ultrafree-MC UFC3 LCC (molecular weight cut-off, 5000; Millipore, Bedford, MA). The filtrate was dried and resolved in 50 µl of pure water. Metabolites were identified and quantified using CE-TOFMS system (Agilent Technologies, Santa Clara, CA).

Enzyme assay. Dehydrogenases for glycerol and glycerol-3-phosphate were assayed according to a previously described method.⁴⁴ Briefly, WT cells were inoculated in 50 ml of fresh S-MB or G-MB, and cultured to the exponential or stationary growth phase. The cells were washed twice with 10 mM potassium phosphate buffer (pH 7.0), and resuspended in 2 ml of the same buffer. The cells were ultrasonically disrupted at 4°C and 9 kHz for 10 min (Insonator model 201M; Kubota, Tokyo, Japan). Insoluble substances were removed by centrifugation (20,000 ×g, 4°C, 15 min), and the supernatant was used as the cell extract for analysis. The cell extract (20 µl) was mixed with 10 mM (final concentration) potassium phosphate buffer (pH 7.0), 5 mM substrate (glycerol or glycerol-3-phosphate), and 0.5 mM coenzyme (NAD⁺ or NADP⁺). The change in the absorbance of the reaction mixture at a wavelength of 340 nm was monitored at 30°C for 10 min.

Quantitative PCR. Total RNA extraction, *in vitro* cDNA synthesis, and quantitative

PCR were performed to monitor the expression levels of specific genes in the bacterial samples. *A. vinelandii* grown in S-MB or G-MB was harvested at the exponential and stationary growth phases. Total RNA was extracted using the standard hot phenol method. DNA degradation and subsequent RNA purification were performed using the RNase-Free DNase Set (Qiagen, Tokyo, Japan) and an RNeasy Mini Kit (Qiagen), respectively. cDNA, synthesized from a 200 ng RNA sample as using the ReverTra Ace qPCR RT Kit (Toyobo, Osaka, Japan) according to the manufacturer's instructions, was used as a template. Quantitative PCR was performed using the SYBR Premix Ex Taq GC (Takara Bio, Shiga, Japan) and the LineGene instrument (Toyobo).

Electron microscopy. TEM and SEM analyses were entrusted to Tokai Electron Eicroscopy Analysis Co. (Nagoya, Japan). *A. vinelandii* was grown in S-MB or G-MB media (30 ml of working volume) for 72 h. In the case of TEM analysis, the culture and fixing solution A (2% paraformaldehyde, 2% glutaraldehyde, and 0.1 M potassium phosphate buffer, pH 7.4) were mixed in a ratio of 1:1 (total 1 ml), and stored at 4°C for 60 min. Cell pellets were collected by centrifugation (5000 ×g, 4°C, 5 min). Fixing solution B (2% osmium tetroxide and 0.1 M potassium phosphate buffer, pH 7.4) was added to each cell pellet and agitated. Preparation of an ultrathin section and TEM analysis using a JEM-1200EX instrument (JEOL, Tokyo, Japan) were carried out as described previously.⁴³ In the case of SEM analysis, the bacterial cells were prefixed at 4°C for 1 h by mixing the culture with an equal volume of fixative consisting of 4% paraformaldehyde and 4% glutaraldehyde in 100 mM cacodylate buffer (pH 7.4). After centrifugation, the bacterial cells were fixed at 4°C for 24 h with 2% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.4), additionally fixed at 4°C for 2 h with 1% tannic acid

in 0.1 M cacodylate buffer (pH 7.4), and washed at 4°C with the same buffer. The fixed cells were treated at 4°C for 3 h with 2% OsO₄ in 0.1 M cacodylate buffer (pH 7.4) and dehydrated in a series of ethanol (50%, 70%, 90%, and 100%, each 30 min). After dehydration, the cells were continuously dehydrated with 100% ethanol at room temperature overnight. The cells were substituted into tert-butyl alcohol at room temperature, followed by freeze drying under vacuum. The cells were coated with a thin layer (60 nm) of osmium by an osmium plasma coater (NL-OPC80NS, Nippon Laser & Electronics Laboratory, Nagoya, Japan). The cells were observed using a scanning electron microscope (JSM-6340F, JEOL, Tokyo, Japan) at an acceleration voltage of 5.0 kV.

PHB detection. Intracellular PHB was detected according to a previously described procedure with slight modification.^{45,46} Dried cells were treated at 100°C for 140 min with 1 ml chloroform containing 0.5% (w/v) benzoic acid as an internal standard and 1 ml methanol containing 3% sulfuric acid. After addition of 1 ml distilled water, each solution was agitated for 1 min and centrifuged (1000 ×g, 4°C, 5 min). The lower organic solvent layer of each sample was subjected to gas chromatography (GC) analysis using a GC-2014 instrument (Shimadzu, Kyoto, Japan) and a DB-5 column (30 m × 0.25 mm × 0.25 μm; Agilent Technologies). Helium was used as the carrier gas. Detector and injector temperatures were set to 275 and 230°C, respectively. Initial and end temperatures were set to 60 and 200°C, respectively, with a gradient of 8°C/min. An authentic sample of PHB (Sigma, St. Louis, MO) was also treated as described above and was used for the identification and quantitative determination of PHB.

Isolation of alginate-overproducing mutant. *A. vinelandii* was grown in G-MB to the exponential growth phase (approximately 0.8 at OD₆₀₀), and the harvested cells obtained by centrifugation were washed with MB buffer (i.e., MB without a carbon source). To create random mutation in the species, the cells were treated with 50 µg/ml *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (NTG; Sigma) at 30°C for 30 min. The NTG solution was removed from the sample by centrifugation, and the cells were incubated overnight in S-MB. After colony formation on an S-MB plate, the highly mucoid cells were selected as candidates for alginate-overproducing mutants.

Alginate assay. Alginate was assayed according to the method of Knutson and Jeanes.⁴⁷ Briefly, the culture broth (200 µl) was mixed with 0.5 M EDTA (12 µl) and 5 M NaCl (4 µl). Cells were removed from the mixture by centrifugation (7000 ×g, room temperature, 5 min). The supernatant (87.5 µl) was mixed with an ice-chilled mixture of sulfuric acid (732.5 µl) and boric acid solution (17.5 µl; 45 mM KOH and 1 M boric acid) as well as 0.1% (w/v) carbazol (25 µl). The mixture was incubated at 55°C for 30 min, and its absorbance 530 nm was subsequently measured. The alginate concentration in the culture broth was determined based on the calibration using seaweed alginate as a standard.

ACKNOWLEDGMENTS

This work was supported by Promotion of Basic Research Activities for Innovative Bioscience (PROBRAIN) in Japan (to K.M.), Grants-in-Aid from the Japan Society for the Promotion of Science (to K.M. and W.H.), and by the Research Grant (to W.H.) from Japan Soap and Detergent Association.

334

335 REFERENCES

- 336 1. Kennedy C, Toukdarian A. Genetics of azotobacters: applications to nitrogen
337 fixation and related aspects of metabolism. Annu Rev Microbiol 1987; 41:
338 227-58.
- 339 2. Setubal JC, dos Santos P, Goldman BS, Hertesvåg H, Espin G, LRubio LM,
340 Valla S, Almeida NF, Balasubramanian D, Cromes L, et al. Genome sequence of
341 *Azotobacter vinelandii*, an obligate aerobe specialized to support diverse
342 anaerobic metabolic processes. J Bacteriol 2009; 191: 4534-45.
- 343 3. Burk D, Lineweaver H. The influence of fixed nitrogen on *Azotobacter*. J
344 Bacteriol 1930; 19:389-414.
- 345 4. Lee CC, Hu Y, Ribbe MW. Unique features of the nitrogenase VFe protein from
346 *Azotobacter vinelandii*. Proc Natl Acad Sci USA 2009; 106: 9209-14.
- 347 5. Oelze J. Respiratory protection of nitrogenase in *Azotobacter* species: is a
348 widely held hypothesis unequivocally supported by experimental evidence?
349 FEMS Microbiol Rev 2000; 24: 321-33.
- 350 6. Galindo E, Peña C, Núñez C, Segura D, Espín G. Molecular and bioengineering
351 strategies to improve alginate and polydihydroxyalkanoate production by
352 *Azotobacter vinelandii*. Microb Cell Fact 2007; 6: 7.
- 353 7. Page WJ, Knosp O. Hyperproduction of poly-β-hydroxybutyrate during
354 exponential growth of *Azotobacter vinelandii* UWD. Appl Environ Microbiol
355 1989; 55: 1334-9.
- 356 8. Remminghorst U, Rehm BH. Bacterial alginates: from biosynthesis to
357 applications. Biotechnol Lett 2006; 28: 1701-12.

- 358 9. Sabra W, Zeng AP, Deckwer WD. Bacterial alginate: physiology, product quality
359 and process aspects. *Appl Microbiol Biotechnol* 2001; 56: 315-25.
- 360 10. Keshavarz T, Roy I. Polyhydroxyalkanoates: bioplastics with a green agenda.
361 *Curr Opin Microbiol* 2010; 13: 321-6.
- 362 11. Braunegg G, Lefebvre G, Genser KF. Polyhydroxyalkanoates, biopolyesters from
363 renewable resources: physiological and engineering aspects. *J Biotechnol* 1998;
364 65:127-61.
- 365 12. Parker LT, Socolofsky MD. Central body of the *Azotobacter* cyst. *J Bacteriol*
366 1966; 91: 297-303.
- 367 13. Pindar DF, Bucke C. The biosynthesis of alginic acid by *Azotobacter vinelandii*.
368 *Biochem J* 1975; 152: 617-22.
- 369 14. Stevenson LH, Socolofsky MD. Cyst formation and poly- β -hydroxybutyric acid
370 accumulation in *Azotobacter*. *J Bacteriol* 1966; 91: 304-10.
- 371 15. Remminghorst U, Rehm BH. In vitro alginate polymerization and the functional
372 role of Alg8 in alginate production by *Pseudomonas aeruginosa*. *Appl Environ*
373 *Microbiol* 2006; 72: 298-305.
- 374 16. Peralta-Gil M, Segura D, Guzmán J, Servín-González L, Espín G. Expression of
375 the *Azotobacter vinelandii* poly- β -hydroxybutyrate biosynthetic *phbBAC* operon
376 is driven by two overlapping promoters and is dependent on the transcriptional
377 activator PhbR. *J Bacteriol* 2002; 184: 5672-7.
- 378 17. Castañeda M, Guzmán J, Moreno S, Espín G. The GacS sensor kinase regulates
379 alginate and poly- β -hydroxybutyrate production in *Azotobacter vinelandii*. *J*
380 *Bacteriol* 2000; 182:2624-8.
- 381 18. Castañeda M, Sánchez J, Moreno S, Núñez C, Espín G. The global regulators

- 382 GacA and σ^S form part of a cascade that controls alginate production in
383 *Azotobacter vinelandii*. J Bacteriol 2001; 183: 6787-93.
- 384 19. Martínez-Salazar JM, Moreno S, Nájera R, Boucher JC, Espín G
385 Soberón-Chávez G, Deretic V. Characterization of the genes coding for the
386 putative sigma factor AlgU and its regulators MucA, MucB, MucC, and MucD
387 in *Azotobacter vinelandii* and evaluation of their roles in alginate biosynthesis. J
388 Bacteriol 1996; 178: 1800-8.
- 389 20. Moreno S, Nájera R, Guzmán J, Soberón-Chávez G, Espín G. Role of alternative
390 sigma factor *algU* in encystment of *Azotobacter vinelandii*. J Bacteriol 1998;
391 180: 2766-9.
- 392 21. Núñez C, León R, Guzmán J, Espín G, Soberón-Chávez, G. Role of *Azotobacter*
393 *vinelandii* *mucA* and *mucC* gene products in alginate production. J Bacteriol
394 2000; 182: 6550-6.
- 395 22. Pyla R, Kim TJ, Silva JL, Jung YS. Overproduction of poly- β -hydroxybutyrate
396 in the *Azotobacter vinelandii* mutant that does not express small RNA ArrF.
397 Appl Microbiol Biotechnol 2009; 84: 717-24.
- 398 23. Segura D, Espín G. Mutational inactivation of a gene homologous to *Escherichia*
399 *coli* *ptsP* affects poly- β -hydroxybutyrate accumulation and nitrogen fixation in
400 *Azotobacter vinelandii*. J Bacteriol 1998; 180: 4790-8.
- 401 24. Senior PJ, Dawes EA. The regulation of poly- β -hydroxybutyrate metabolism in
402 *Azotobacter beijerinckii*. Biochem J 1973; 134: 225-38.
- 403 25. Wu G, Moir AJ, Sawers G, Hill S, Poole RK. Biosynthesis of
404 poly- β -hydroxybutyrate (PHB) is controlled by CydR (Fnr) in the obligate
405 aerobe *Azotobacter vinelandii*. FEMS Microbiol Lett 2001; 194:215-20.

- 406 26. Horan NJ, Jarman TR, Dawes EA. Effects of carbon source and inorganic
407 phosphate concentration on the production of alginic acid by a mutant of
408 *Azotobacter vinelandii* and on the enzymes involved in its biosynthesis. J Gen
409 Microbiol 1981; 127: 185-91.
- 410 27. Page WJ, von Tigerstrom M. Induction of transformation competence in
411 *Azotobacter vinelandii* iron-limited cultures. Can J Microbiol 1978; 24: 1590-4.
- 412 28. Coombs A. Glycerin bioprocessing goes green. Nat Biotechnol 2007; 25: 953-4.
- 413 29. da Silva GP, Mack M, Contiero J. Glycerol: a promising and abundant carbon
414 source for industrial microbiology. Biotechnol Adv 2009; 27: 30-9.
- 415 30. Bizzini A, Zhao C, Budin-Verneuil A, Sauvageot N, Giard JC, Auffray Y, Hartke
416 A. Glycerol is metabolized in a complex and strain-dependent manner in
417 *Enterococcus faecalis*. J Bacteriol 2010; 192: 779-85.
- 418 31. Wang ZX, Zhuge J, Fang H, Prior BA. Glycerol production by microbial
419 fermentation: a review. Biotechnol Adv 2001; 19: 201-23.
- 420 32. Wendisch VF, Bott M, Eikmanns BJ. Metabolic engineering of *Escherichia coli*
421 and *Corynebacterium glutamicum* for biotechnological production of organic
422 acids and amino acids. Curr Opin Microbiol 2006; 9: 268-74.
- 423 33. Petty F. GABA and mood disorders: a brief review and hypothesis. J Affect
424 Disord 1995; 34: 275-81.
- 425 34. Castanie-Cornet MP, Penfound TA, Smith D, Elliott JF, Foster JW. Control of
426 acid resistance in *Escherichia coli*. J Bacteriol 1999; 181: 3525-35.
- 427 35. Page WJ, Manchak J, Rudy B. Formation of
428 poly(hydroxybutyrate-co-hydroxyvalerate) by *Azotobacter vinelandii* UWD.
429 Appl Environ Microbiol 1992; 58: 2866-73.

- 430 36. Zhu C, Nomura CT, Perrotta JA, Stipanovic AJ, Nakas JP. Production and
431 characterization of poly-3-hydroxybutyrate from biodiesel-glycerol
432 by *Burkholderia cepacia* ATCC 17759. Biotechnol Prog 2010; 26: 424-30.
- 433 37. Cui B, Huang S, Xu F, Zhang R, Zhang Y. Improved productivity of poly
434 (3-hydroxybutyrate) (PHB) in thermophilic *Chelatococcus daeguensis* TAD1
435 using glycerol as the growth substrate in a fed-batch culture. Appl Microbiol
436 Biotechnol 2015; In press.
- 437 38. Cavaleiro JMBT, de Almeida MCMD, Grandfils C, da Fonseca
438 MMR. Poly(3-hydroxybutyrate) production by *Cupriavidus necator* using waste
439 glycerol. Process Biochem 2009; 44: 509-15.
- 440 39. Tanadchangsang N, Yu J. Microbial synthesis of polyhydroxybutyrate from
441 glycerol: gluconeogenesis, molecular weight and material properties of
442 biopolyester. Biotechnol Bioeng 2012; 109:2808-18.
- 443 40. Kalaiyezhini D, Ramachandran KB. Biosynthesis of Poly-3-Hydroxybutyrate
444 (PHB) from Glycerol by *Paracoccus denitrificans* in a Batch Bioreactor: Effect
445 of Process Variables. Prep Biochem Biotechnol 2015; 45:69-83.
- 446 41. Ashby R, Solaiman D, Strahan G. Efficient utilization of crude glycerol as
447 fermentation substrate in the synthesis of poly(3-hydroxybutyrate)
448 biopolymers. J Am Oil Chem Soc 2011; 88: 949-59.
- 449 42. Ibrahim MHA, Steinbüchel A. Poly(3-hydroxybutyrate) production from
450 glycerol by *Zobellella denitrificans* MW1 via high-cell-density fed-batch
451 fermentation and simplified solvent extraction. Appl Environ
452 Microbiol 2009; 75: 6222-31.
- 453 43. Hashimoto W, Miyamoto Y, Yamamoto M, Yoneyama F, Murata K. A novel

- 454 bleb-dependent polysaccharide export system in the nitrogen-fixing *Azotobacter*
455 *vinelandii* subjected to low nitrogen gas levels. Int Microbiol 2013; 16: 35-44.
- 456 44. Matsuzawa T, Ohashi T, Hosomi A, Tanaka N, Tohda H, Takegawa K. The *gldI*⁺
457 gene encoding glycerol dehydrogenase is required for glycerol metabolism in
458 *Schizosaccharomyces pombe*. Appl Microbiol Biotechnol 2010; 87: 715-27.
- 459 45. Braunegg G, Sonnleitner B, Lafferty RM. A rapid gas chromatographic method
460 for the determination of poly-3-hydroxybutyric acid in microbial biomass. Eur J
461 Appl Microbiol Biotechnol 1978; 6: 29-37.
- 462 46. Tsuge T, Tanaka K, Ishizaki A. Development of a novel method for feeding a
463 mixture of L-lactic acid and acetic acid in fed-batch culture of *Ralstonia*
464 *eutropha* for poly-D-3-hydroxybutyrate production. J Biosci Bioeng 2001; 91:
465 545-50.
- 466 47. Knutson CA, Jeanes A. Determination of the composition of uronic acid
467 mixtures. Anal Biochem 1968; 24: 482-90.
468

FIGURE LEGENDS

Figure 1. Putative synthetic pathway for PHB and alginate from glycerol in *A. vinelandii*. The genes involved in glycerol metabolism are also indicated.

Figure 2. Gene expression of glycerol-3-phosphate and glycerol dehydrogenases. (A) Total RNA from *A. vinelandii* grown in G-MB or S-MB to the exponential and stationary growth phases. (B) Relative gene expression of glycerol-3-phosphate (left) and glycerol dehydrogenases (right). For quantitative PCR, cDNA synthesized from was used as a template in all samples. The expression levels of both genes in S-MB at the stationary phase were standardized as the relative expression of 1.

Figure 3. PHB granules in *A. vinelandii* cells revealed by TEM. Bacteria were cultured for 72 h in S-MB or G-MB as indicated.

Figure 4. Time course of PHB production by *A. vinelandii*. (A) Cell growth. (B) PHB accumulation in dry cells. Open circles, WT grown on sucrose; closed circles, WT grown on glycerol; open triangles, $\Delta algD$ grown on sucrose; closed triangles, $\Delta algD$ grown on glycerol.

Figure 5. Alginate-overproducing mutants of *A. vinelandii*. (A) TEM observation of MT1 cells grown on glycerol. (B) TEM observation of WT cells grown on glycerol. (C) SEM observation of WT cells grown on glycerol. Panels A' B', and C' are magnified views of the regions of interest indicated in panels A, B, and C, respectively.

- 1 **Figure 6.** Alginate production from glycerol-grown *A. vinelandii*. (A) Cell growth. (B)
- 2 Extracellular alginate. Circles, WT; triangles, alginate-overproducing mutant.
- 3

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.





Figure 6.



1 Table 1. Intracellular metabolites determined through metabolomics.

Metabolite	Concentration (pmol/OD ₆₀₀ ml)			
	Sucrose		Glycerol	
	Log phase	Stationary phase	Log phase	Stationary phase
Glucose 6-phosphate ^{*1}	18.7 (±4.2)	22.5 (±29.3)	22.6 (±5.2)	61.0 (±25.4)
6-Phosphogluconic acid	U. L.	U. L.	U. L.	U. L.
2-Keto-3-deoxy-6-phosphogluconic acid	N. D.	N. D.	N. D.	N. D.
Fructose 6-phosphate ^{*2}	8.3 (±1.7)	8.2 (±11.6)	13.9 (±6.3)	29.4 (±13.1)
Mannose 6-phosphate ^{*1}	18.7 (±4.2)	22.5 (±29.3)	22.6 (±5.2)	61.0 (±25.4)
Mannose 1-phosphate ^{*2}	8.3 (±1.7)	8.2 (±11.6)	13.9 (±6.3)	29.4 (±13.1)
GDP Mannose	0.3 (±0.1)	1.0 (±0.2)	2.1 (±0.6)	5.5 (±1.3)
GDP Mannuroic acid	N. D.	N. D.	N. D.	N. D.
Glycerol-3-phosphate	11.0 (±4.4)	4.6 (±3.1)	400.3 (±186.4)	82.0 (±61.9)
Dihydroxyacetone phosphate	0.8 (±0.2)	2.0 (±1.9)	12.6 (±5.3)	8.5 (±2.5)
Fructose 1,6-biphosphate	0.5 (±0.1)	1.3 (±0.1)		2.3 (±0.4)
Glyceraldehyde-3-phosphate	2.4 (±0.4)	1.7 (±1.0)	6.2 (±2.0)	7.6 (±0.6)
Glycerate 1,3-biphosphate	N. D.	N. D.	N. D.	N. D.
3-Phosphoglyceric acid	0.8 (±0.2)	5.0 (±4.1)	1.4 (±0.1)	10.3 (±5.5)
2-Phosphoglyceric acid	N. D.	N. D.	N. D.	N. D.
Phosphoenolpyruvic acid	U. L.	7.3 (±0.1)	U. L.	14.4 (±4.0)
Pyruvic acid	11.5 (±3.7)	9.7 (±3.7)	42.5 (±24.8)	12.7 (±14.2)
Acetyl CoA	2.5 (±1.5)	4.8 (±3.8)	2.3 (±0.2)	9.1 (±4.6)

2 U. L., undetectable level; N. D., not detectable under this experimental condition. ^{*1},
3 summation of glucose 6-phosphate and mannose 6-phosphate; ^{*2}, summation of fructose
4 6-phosphate and mannose 1-phosphate. Data are averages (± SD) of three experiments.

5

1 Table 2. Intracellular amino acids and other organic acids.

Metabolite	Concentration (pmol/OD ₆₀₀ ml)			
	Sucrose		Glycerol	
	Log phase	Stationary phase	Log phase	Stationary phase
Ala	121.9 (±18.1)	95.0 (±61.3)	220.9 (±88.1)	114.1 (±35.9)
Arg	55.3 (±13.4)	19.7 (±26.0)	51.6 (±26.0)	93.7 (±16.4)
Asn	1.7 (±1.1)	U. L.	1.9 (±1.4)	5.9 (±7.4)
Asp	14.2 (±4.1)	4.9 (±3.8)	24.5 (±4.1)	24.3 (±28.1)
Cys	U. L.	U. L.	U. L.	U. L.
Gln	134.3 (±8.8)	146.3 (±180.5)	217.1 (±158.1)	184.1 (±73.3)
Glu	85.7 (±25.4)	75.4 (±112.6)	368.6 (±127.6)	134.3 (±106.9)
Gly	45.5 (±22.1)	9.5 (±5.4)	65.1 (±9.3)	70.3 (±84.6)
His	4.4 (±1.6)	1.4 (±0.8)	4.3 (±1.9)	13.3 (±12.8)
Ile	18.8 (±5.1)	1.1 (±0.5)	30.3 (±13.3)	8.9 (±11.7)
Leu	45.1 (±11.5)	1.7 (±0.7)	76.8 (±45.6)	13.0 (±17.3)
Lys	47.4 (±7.6)	13.1 (±12.9)	69.1 (±21.1)	83.5 (±32.1)
Met	7.8 (±1.8)	0.3 (±0.1)	8.5 (±5.9)	2.3 (±2.4)
Phe	7.5 (±2.9)	1.0 (±0.3)	14.9 (±8.1)	7.8 (±7.4)
Pro	15.4 (±3.3)	4.1 (±4.2)	28.2 (±5.7)	31.3 (±12.8)
Ser	58.5 (±36.7)	14.3 (±9.7)	68.1 (±23.7)	115.2 (±137.9)
Thr	39.3 (±9.8)	4.3 (±3.1)	65.0 (±25.8)	34.9 (±29.8)
Trp	2.0 (±1.0)	0.4 (±0.3)	2.0 (±1.0)	0.8 (±0.8)
Tyr	22.0 (±5.2)	3.5 (±3.2)	44.3 (±29.7)	20.7 (±10.3)
Val	121.6 (±37.6)	3.4 (±1.6)	168.8 (±100.8)	22.8 (±24.4)
γ-Aminobutyric acid (GABA)	113.9 (±39.6)	0.8 (±0.2)	238.9 (±278.8)	1.7 (±2.1)
Lactic acid	501.0 (±61.1)	530.5 (±309.1)	1239.3 (±507.3)	719.2 (±646.8)

2 U. L., undetectable level. Data are averages (± SD) of three experiments.

3

1 Table 3. Bacterial strains used in the present study.

Bacterial strains (abbreviation)	Characteristics or sequences	References
<i>Azotobacter vinelandii</i> ATCC 12837 (WT)	Wild type strain	ATCC
<i>A. vinelandii</i> $\Delta algD$ ($\Delta algD$)	Alginate-deficient strain with an insertion of the tetracycline resistance gene in chromosomal <i>algD</i> .	43
<i>A. vinelandii</i> MT1 (MT1)	Alginate-overproducing strain derivatized from WT by NTG treatment.	Present study

2 ATCC, American Type Culture Collection.